arXiv:1201.2440v1 [cond-mat.quant-gas] 11 Jan 2012

Self-consistent field theory of polarized BEC: dispersion of collective excitation

P. A. Andreev* and L. S. Kuzmenkov†
Faculty of Physics, Moscow State University, Moscow, Russian Federation.

(Dated: January 13, 2012)

We propose the construction of a set of quantum hydrodynamics equations for the Bose-Einstein condensate (BEC) where atoms have electric dipole moment. The contribution of the dipole-dipole interactions (DDI) to the Euler equation is estimated. Quantum equations for the evolution of medium polarization are constructed. The mathematical method we developed allows studying the effects of interactions on the evolution of polarization. The developed method may be applied to various physical systems in which dynamics is affected by DDI. Derivation of Gross-Pitaevskii equation for polarized particles is discussed using point of view of quantum hydrodynamics. We shown that Gross-Pitaevskii equation appear at condition there all dipoles has the same direction which does not change in time. Comparison of equation of electric dipole interaction with equation of magnetization evolution is described. A problem of elementary excitations in BEC, either affected or not affected by the uniform external electric field, is considered using our method. We show that the evolution of polarization in BEC leads to the formation of a novel type of elementary excitations. Detailed description of dispersion of elementary excitations is presented. We consider also the process of wave generation in polarized BEC by means of monoenergetic beam of neutral polarized particles. We compute the possibilities of the generation of Bogoliubov's modes and polarization modes by the dipole beam.

I. INTRODUCTION

After obtaining the Bose-Einstein condensate (BEC) in experiment with pairs of alkaline metal atoms, theoretical and experimental investigation of linear waves and nonlinear structures in it have been performed. In recent years the interest to polarized BEC has been increasing. It connected with recent experimental progress in cooling of polarized atoms and molecules. At present day the BEC with magnetic polarization is realized on atoms ^{52}Cr. There are a lot of attempt of experimental obtaining of electrically polarized BEC (see review [1]- [4]). For this aim BEC of molecules has been cooled. Particular interest to electrically polarized BEC brought because of large dipole-dipole scattering length, and thereby because of both the big magnitude and large distance of interaction compare with analogous quantities of the magnetized BEC.

Many processes in quantum systems are determined by the dynamics and dispersion of elementary excitations (EE) [5]. A law of dispersion of EE in degenerate dilute Bose gas was obtained by Bogoliubov in 1947 [6, 7]. Many authors studied the change of the Bogoliubov's mode which arises when SRI interactions accounting more carefully [8]- [14], or geometry of the system is complex [15], [16], [17]. In articles [18]- [22] authors studied the influence of electric dipole moment (EDM) dynamics on dispersion of EE in BEC. The contribution of polarization in dispersion law of Bogoliubov's mode was obtained in [18]- [22]. Instability of the Bogoliubov's mode in 3D dipoles BEC (DBEC) with repulsive short range interaction (SRI) were shown in Ref. [18]- [20]. U. R. Fisher [21] obtained that Bogoliubov's mode in 2D DBEC is stable for a wide range of system parameters. There are also review papers [2]- [4] where various aspects of physics of polarized BEC were considered. In this paper we interested in possibilities of polarization waves existence in DBEC, i.e. another type of EE. Dynamics of polarization is interesting not for quantum gases only, but in solid state physics and physics of low-dimensional systems too. The polarization waves in low dimensional and multy-layer systems of conductors and dielectrics are considered in the papers [23, 24]. Analogously, in BEC of atoms with EDM we expect the existence of polarizations wave; along with Bogoliubov's mode.

For investigation of BEC, in system of particles with EDM or magnetic moments, various theoretical methods are used. One of the ways of theoretical description of the polarized BEC is the generalization of well-known Gross-Pitaevskii (GP) equation [25]- [29]. In this way next papers were made: for the spinor BEC [30–32], for the effect of magnetic moment on the BEC evolution [32]- [41] and for the influence of electrical polarization of atoms on the dynamic processes in BEC [32]- [42]. Dipolar fermions also attract a lot of attention [43], [44].

The same generalization of GP equation usually used for magnetic moments and electric dipoles:

$$\imath\hbar\partial_t\Phi(\mathbf{r},t)=\biggl(-\frac{\hbar^2\nabla^2}{2m}+\mu(\mathbf{r},t)+V_{ext}(\mathbf{r},t)+g\mid\Phi(\mathbf{r},t)\mid^2$$

$$+d^2\int d\mathbf{r}'\frac{1-3\cos^2\theta'}{|\mathbf{r}-\mathbf{r}'|^3}\mid\Phi(\mathbf{r}',t)\mid^2\biggr)\Phi(\mathbf{r},t). \qquad (1)$$

In equation (1) following designation are used: $\Phi(\mathbf{r},t)$ is the macroscopic wave function, μ is the chemical potential, V_{ext} is the potential of external field, g is the constant of short-range interaction, d is the dipole electric moment of single atom, m is the mass of particles and $\hbar$ is the Planck constant divided by 2π.

Along with GP equation the method of quantum hydrodynamics (QHD) is also used. The method of QHD [14],

[45], [46] was used for investigations of various physical systems, particularly for BEC [14], [47] and polarization waves in conductors and semiconductors [23]. Therefore, the method of QHD could be used to consider the possibility of arising polarization waves in BEC and calculating of the dispersion law of these waves.

At interpretation of DDI in quantum gases (QG) used the notion of scattering length (SL) and first Born approximation (FBA), analogously to the SRI described by the fourth term in right-side of equation (1). Therefore, the problem reduces to the process of scattering. Different approximations based on scattering process was analyzed in Ref. [48]. Particularly, where were considered condition there can be used FBA and presented generalization of GP equation for the scattering of polarized atoms beyond FBA.

There we consider QHD we do not formulate the scattering problem. We do not need to confine ourselves to a scattering process, because we interesting in interaction between atoms including interaction between several atoms at the same time. This statement is especially important for polarized atoms because of the long-range interaction between dipoles. Of course for system of particles with low concentration each act of interaction between dipoles could be considered as a scattering process. This approximation is possible as for neutral polarized particles as for charged particles. The kinetic theory for the last case was developed by L. D. Landau in 1936 [49]. Landau constructed collision term for Coulomb particles, using analogy with the Boltzmann equation. Landau collision integral depend on scattering cross section. In this connection he considered the problem of scattering of charged particles on small angle, that is suitable for rare systems. Strictly speaking, for particles with long-range interaction the conception of self-consistent field is more suitable. This conception was suggested by A. A. Vlasov in 1938, for system of charged particles [50]. In fact group of particles moves in the field of their neighbors, but state of motion of neighbors depend on motion of this group of particles as well (This picture of interaction is more detailed described on Fig. 1). According to this idea Vlasov proposed kinetic equation for charged particles called Vlasov's equation. Actually the Landau collision integral and the Vlasov's equation are the two opposite branches of hypothetical general theory. In this paper we follow to Vlasov's approximation and derive the self-consistent field theory for dipole-dipole interaction in quantum gases. From this point of view there no need to consider the first Born approximation or scattering process as well. In Appendix B to this paper we present derivation of GP equation for polarized particles from QHD equations. We did it at the condition that dipoles are parallel each other and to external field, they directions does not oscillate or vibrate around equilibrium direction.

Other methods were also applied for investigation of the polarized BEC and other systems of particles, with EDM, along with GP equation. A hydrodynamic formulation of the Hartree-Fock theory for particles with significant EDM is considered in [51]. In paper [51] the Euler-type equation was derived, from the evolution of the density matrix. The EDM dynamics in dimer Mott insulators causes the rise of the low-frequency mode [52]. A two-body quantum problem for polarized molecules is analyzed in [53]. The existence of states with spontaneous interlayer coherence has been predicted in [54] in systems of polar molecules. Calculations, in [54], were based upon the secondary quantization approach, where the Hamiltonian accounts for the molecular rotation and dipole-dipole interactions. Superfluidity anisotropy of polarized fermion systems was shown in [55] and their thermodynamic and correlation properties were investigated [56]. The effect of EDM in a system of cooled neutral atoms which are used for quantum computing and quantum memory devices was analyzed in [57].



FIG. 1. (Color online) The figure presents the picture of self-consistent interaction between dipoles. Total polarization of region 2 (red) interacts with total polarization of region 1 (green). Changing the extreme point of a radius vector (or shifting the region 2) we can scan whole space. In this way we obtain action of external dipoles on region 1. Changing position of region 1 and repeating described operation we obtain action surrounding dipoles on each region of space. This is a picture of self-consistent interaction in fixed moment of time and this picture governs an evolution of polarization in system. This picture of interaction is typical for classic physics, where we need to obtain smooth functions describing collective motion. For that is necessary to average at physically infinitesimal volume (sketched circle). In quantum mechanics, where concentration, polarization, etc, defined via wave function and we can consider described picture on interaction of separate particle instead of space regions.

The characteristic property of BEC in a system of excitons inducing in semiconductors [58] it is a significant value of EDM of excitons. This leads to strong interaction of excitons with external electric field and emergence of the collective DDI in exciton systems. QHD method may be applied to such systems along with quantum kinetics based on the nonequilibrium Green functions [59] or density matrix equations [60].

Notable success has been reached in the Bose condensation of dense gases [61, 62]. Consequently, we need the



FIG. 2. (Color online) The figure shows motion of dipoles without changing of direction of dipoles during motion. Dipoles stand at temperature equal to zero $T = 0$ and in external uniform static electric field **E**. At lower picture describe particles in two different moments of time. White dipoles presents system in the first moment of time when perturb only one dipole, but violet dipoles are present system in the next moment.



FIG. 3. (Color online) The figure presents the change of polarization density at movement of dipoles in the absence of dipoles direction evolution.

detailed account of short range interaction in investigations of dynamics of EDM. In this work we account the SRI up to the third order on interaction radius (TOIR) [14]. This approximation leads to nonlocality of SRI [14], [63]- [66].

Electrically polarized BEC can interact with the beam of charged and polarized particles by means of charge-dipole and dipole-dipole interaction. Such interaction leads to transfer of energy from beam to medium and, consequently, to generation of waves. In plasma physics the effect of generation of waves by electron [67] or magnetized neutron [68, 69] beam are well-known. When we use term beam we also mean stream of particles excited in the considered system, along with external beam of particles passing through the system. In presented paper we consider similar effect in polarized BEC.

Our treatment we derived and used the QHD equations for polarized particles. System of QHD equations consists of continuity equation, momentum balance equation, equation of polarization evolution and equation of polarization current evolution. We made first principles derivation of this equation. For this purpose we used many-particle Schrödinger equation. In this article we analytically calculate the dispersion properties of polarized BEC. We have shown that the dynamics of EDM leads to existence of new branch in dispersion law. Consequently, there are the waves of polarization in polarized BEC, along with Bogoliubov's mode. We obtain appropriate contribution to the dispersion of Bogoliubov's mode related to DDI. Then, we consider the process of wave generation in polarized BEC by means of monoenergetic beam of neutral polarized particles. We predict new method of generation of Bogoliubov's mode and a new type of modes, rest of the polarization waves.

This paper is the extension and result of processing of our previous paper [70], where the brief description of the same results and ideas were presented.

This paper is organized as follows. We introduce the model Hamiltonian in Sec. II, and present the momentum balance equation. Farther, in Sec. II, we derive the equation of evolution of polarization and obtain the influence of the interactions on the evolution of polarization. In Sec. III we calculate the dispersion dependence of EE in BEC. The polarization evolution is taken into account. We obtain the contribution of polarization in dispersion of Bogoliubov's mode and show the existence of new wave solution. We consider high frequency excitation appearing at large equilibrium polarizations. We also present calculations for the wave of polarization at constant concentration. In Sec. IV we study the wave generation in polarized BEC by means of neutral polarized particle beam. In Sec. V we present the brief summary of our results. App. A. contains detail of derivation of equations of polarization evolution. In App. B. derivation of GP equation for polarized particles is presented. In App. C. we consider distinction of evolution of spinning particle from the particles with EDM.

II. THE MODEL

In GP equation term described DDI, for both EDM and magnetic moment, interpret via dipole scattering length (DSL) a_{dd}. The angle between dipoles coming in GP equation via potential of DDI, and equation is solved for various angles between dipoles. But equation does not contain information about evolution of dipoles direction or, for the case of soft dipoles than dipoles caused by external field, variation of dipole magnitude. Change of dipoles direction there is due to interaction. If we have external static uniform electric field, at close to zero temperatures, all dipoles will be direct parallel to external field. At low temperatures there is no temperature disorientation of dipoles and we have system of parallel dipoles, as it is shown on Fig. 2. In such systems there are two type of dipoles motion. The first one is the motion there the direction of dipoles is preserves Fig. 2, 3. In this case DDI has influence on dipoles properties because it is change the strength of interaction and leads to anisotropy of interaction. At such



FIG. 4. (Color online) The figure describes how the polarization direction perturbation propagates through the system of particles. We consider system of dipoles at $T = 0$ and being in external uniform electric field. Upper picture show that if at the first moment there is perturbation of the EDM in next moment this perturbation pass to several neighbors. At lower picture describe particles in two different moments of time. White dipoles presents system in the first moment of time when perturb only one dipole, but violet dipoles are present system in the next moment.

form of motion the polarization is changed accordingly to concentration $\mathbf{P}(\mathbf{r}, t) = \mathbf{d}n(\mathbf{r}, t)$, where $\mathbf{d}$ is the EDM of single particle Fig. 3. The second type of motion is a motion of dipoles including change of dipoles direction Fig.s 4, 5. Perturbation of dipoles direction can be caused by both external influence and local fluctuation of dipoles direction produced by DDI. In consequence of DDI the perturbation will propagate through system. Particularly, this propagation might be the wave nature. For description of such motion we need to know how polarization of system evolves in different point of space during the time. Therefore we must derive equation describing the evolution of polarization field and accounting influence of interaction of this evolution.

Angle dependence in GP equation is important for the case when external electric field is nonuniform. In this case dipoles in different areas of space have different directions Fig. 6 and it is reflected on interaction via angle dependence under integral in GP equation (1). Still, this equation does not contain information about propagation of dipoles direction perturbation.



FIG. 5. (Color online) The figure presents the oscillation of EDM near of the equilibrium position which defined by external electric field **E**. Indexes 1-6 present the different moments of time $t_1 < t_2 < ... < t_6$. This figure describes mechanism of the polarization change due to oscillations of EDM of each particle in the system. This mechanism might reveal along with rotation of EDM. Incoherent motion of EDMs which is combination of rotation and oscillation is presented on Fig. 4(b).



FIG. 6. (Color online) The figure presents the two dipoles with the EDM turned in different directions. At propagation of perturbation through the system relative directions of dipoles is changed. This lied to change of force of interaction in concordance with potential of DDI (2). In Appendix B we shown that GP equation describe motion for parallel dipoles only.

For studying we pay attention for the SRI and DDI. We also consider the action of external fields on particles. The action of external electric field on dipoles we consider in evident form and the external field described trap presented in general form by trapping potential $V_{trap}(\mathbf{r}_i, t)$.

Here we used quasi-static approximation for DDI. The meaning of quasi-static approximation is that we consider interaction between dipoles at fixed moment of time as they are motionless. DDI leads to change of state of motion, and, in particularly, to variation of dipoles direction. Then we talk about change direction we mean both the variation during precession around of rotation axis and deviation of rotation axis direction. Described approximation is an analog of Coulomb interaction there interaction between moving charged approximately described by means of interaction potential of motionless charges, this approximation usually used for description of non-relativistic plasma [73].

Explicit form of the Hamiltonian of considering system in a quasi-static approximation is

$$\hat{H}=\sum_i\left(\frac{1}{2m_i}\hat{\mathbf{p}}_i^2-d_i^\alpha E^\alpha_{i,ext}+V_{trap}(\mathbf{r}_i,t)\right)$$

$$+\frac{1}{2}\sum_{i,j\neq i}\left(U_{ij}-d_i^\alpha d_j^\beta G_{ij}^{\alpha\beta}\right). \tag{2}$$

The first term here is the operator for kinetic energy. The second term represents the interaction between the dipole moment d_i^α and the external electrical field. The subsequent terms represent short-range U_{ij} and dipole-dipole interactions between particles, respectively. The Green's function for dipole-dipole interaction is taken as $G_{ij}^{\alpha\beta}=\nabla_i^\alpha\nabla_i^\beta(1/r_{ij})$.

For investigation of EE dynamic in polarized BEC we derive the system of QHD equations. This system of equations consists of the continuity equation, the Euler's equation and for the case of polarized particles the equations of polarization evolution and equation of field. The system of equations is derived by methods described in [14].

The first equation of a QHD equations system is the continuity equation

$$\partial_t n(\mathbf{r},t)+\partial^\alpha(n(\mathbf{r},t)v^\alpha(\mathbf{r},t))=0. \tag{3}$$

The momentum balance equation for the polarized BEC has the form

$$mn(\mathbf{r},t)(\partial_t+\mathbf{v}\nabla)v^\alpha(\mathbf{r},t)+\partial_\beta p^{\alpha\beta}(\mathbf{r},t)$$

$$-\frac{\hbar^2}{4m}\partial^\alpha\triangle n(\mathbf{r},t)+\frac{\hbar^2}{4m}\partial^\beta\left(\frac{\partial^\alpha n(\mathbf{r},t)\cdot\partial^\beta n(\mathbf{r},t)}{n(\mathbf{r},t)}\right)$$

$$=\Upsilon n(\mathbf{r},t)\partial^\alpha n(\mathbf{r},t)+\frac{1}{2}\Upsilon_2\partial^\alpha\triangle n^2(\mathbf{r},t)$$

$$+P^\beta(\mathbf{r},t)\partial^\alpha E^\beta(\mathbf{r},t), \tag{4}$$

where

$$\Upsilon=\frac{4\pi}{3}\int dr(r)^3\frac{\partial U(r)}{\partial r}, \tag{5}$$

and

$$\Upsilon_2\equiv\frac{\pi}{30}\int dr(r)^5\frac{\partial U(r)}{\partial r}. \tag{6}$$

In equation (4) we defined a parameter Υ_2 as (6). This definition differs from the one in the work [14]. Terms proportional to $\hbar^2$ appear as a result of usage of quantum kinematics. The first two members at the right side of the equation (4) are first terms of expansion of the quantum stress tensor. They occur because of taking into account of the SRI potential U_{ij}. The interaction potential U_{ij} determinates the macroscopic interaction constants Υ and Υ_2. The last two members of the equation (4) describe force fields that affect the dipole moment in a unit of volume as the effect of the external electrical field and the field produced by other dipoles, respectively. The last member is written using the self-consistent field approximation [45], [23]. $p^{\alpha\beta}(\mathbf{r},t)$ is a tensor of the kinetic pressure, which depends on particles' thermal velocities and does not contribute into the BEC dynamics at temperatures near zero.

The first order of the interaction radius interaction constant for dilute gases has the form

$$\Upsilon=-\frac{4\pi\hbar^2a}{m}, \tag{7}$$

where a is the scattering length SL [7, 14]. The value Υ_2 may be expressed approximately as

$$\Upsilon_2=-\frac{\theta a^2\Upsilon}{8}, \tag{8}$$

where θ is a constant positive value about 1, which depends on the interatomic interaction potential. Finally, Υ_2 takes the form

$$\Upsilon_2=-\frac{\pi\theta\hbar^2a^3}{2m}. \tag{9}$$

We have also a field equation

$$\nabla\mathbf{E}(\mathbf{r},t)=-4\pi\nabla\mathbf{P}(\mathbf{r},t). \tag{10}$$

In the case particles does not contain the dipole moment, the continuity equation and the momentum balance equation form a closed system of equations. When the dipole moment is taken into account in a momentum balance equation, a new physical value emerges, a polarization vector field $P^\alpha(\mathbf{r},t)$. This causes system of equations to become incomplete.

Next equation we need for investigation of EE dispersion is the equation of polarization evolution

$$\partial_t P^\alpha(\mathbf{r},t)+\partial^\beta R^{\alpha\beta}(\mathbf{r},t)=0, \tag{11}$$

$R^{\alpha\beta}(\mathbf{r},t)$ is the current of polarization.

The equation (11) does not contain information about the effect of the interaction on the polarization evolution. The evolution equation of $R^{\alpha\beta}(\mathbf{r},t)$ can be constructed by analogy with the above derived equations. Method of the equations derivation is described in Appendix A. Using a self-consistent field approximation of the dipole-dipole interaction we obtain an equation for the polarization current $R^{\alpha\beta}(\mathbf{r},t)$ evolution

$$\partial_t R^{\alpha\beta}(\mathbf{r},t)+\partial^\gamma\biggl(R^{\alpha\beta}(\mathbf{r},t)v^\gamma(\mathbf{r},t)$$

$$+R^{\alpha\gamma}(\mathbf{r},t)v^\beta(\mathbf{r},t)-P^\alpha(\mathbf{r},t)v^\beta(\mathbf{r},t)v^\gamma(\mathbf{r},t)\biggr)$$

$$+\frac{1}{m}\partial^\gamma r^{\alpha\beta\gamma}(\mathbf{r},t)-\frac{\hbar^2}{4m^2}\partial_\beta\triangle P^\alpha(\mathbf{r},t)$$

$$+\frac{\hbar^2}{8m^2}\partial^\gamma\left(\frac{\partial_\beta P^\alpha(\mathbf{r},t)\partial_\gamma n(\mathbf{r},t)}{n(\mathbf{r},t)}+\frac{\partial_\gamma P^\alpha(\mathbf{r},t)\partial_\beta n(\mathbf{r},t)}{n(\mathbf{r},t)}\right)$$

$$= \frac{1}{m}\Upsilon\partial^{\beta}\Bigl(n(\mathbf{r},t)P^{\alpha}(\mathbf{r},t)\Bigr)$$

$$+\frac{\sigma}{m}\frac{P^{\alpha}(\mathbf{r},t)P^{\gamma}(\mathbf{r},t)}{n(\mathbf{r},t)}\partial^{\beta}E^{\gamma}(\mathbf{r},t). \qquad (12)$$

Here $r^{\alpha\beta\gamma}(\mathbf{r},t)$ represents the contribution of thermal movement of polarized particles into the dynamics of $R^{\alpha\beta}(\mathbf{r},t)$. As we deal with BEC below, the contribution of $r^{\alpha\beta\gamma}(\mathbf{r},t)$ may be neglected. The last term of the formula (12) includes both external electrical field and a self-consistent field that particle dipoles create. This term contain numerical constant σ. Used in formula (12) approximations are described in Appendix A.

The first term in right side of Eq. (12) describe the short range interaction.

We can see various interactions are included in the equations (4) and (12) additively. At a short distances among particles acts both SRI and dipole-dipole interaction. At large distances remain only dipole-dipole interaction.

From Eq. (12) we can see that the change of polarization arise from both the dipole-dipole interaction and the short range interaction. The SRI among particle leads to displacement of particles. Consequently, as particles has EDM, where are motion of EDM, e.i. changing of the $R^{\alpha\beta}(\mathbf{r},t)$.

The terms that are proportional to $\hbar^2$ are of quantum origin as they are analogs of the Bohm quantum potential in the momentum balance equilibrium.

In the Appendix C we consider the difference between equations of electrical and magnetic polarization evolution.

QHD of spinning particles with spin-orbit interaction were developed in Ref. [69] for system of charged particles, however where obtained field of the force can be used for neutral quantum gases too.

III. ELEMENTARY EXCITATIONS IN THE POLARIZED BEC

We can analyze the linear dynamics of elemental excitations in the polarized three dimensional BEC using the QHD equations (3), (4), (10), (11) and (12). Let's assume the system is placed in an external electrical field $\mathbf{E}_0 = E_0\mathbf{e}_z$. The values of concentration n_0 and polarization $\mathbf{P}_0 = \kappa\mathbf{E}_0$ for the system in an equilibrium state are constant and uniform and its velocity field $v^{\alpha}(\mathbf{r},t)$ and tensor $R^{\alpha\beta}(\mathbf{r},t)$ values are zero.

We consider the small perturbation of equilibrium state like

$$n = n_0 + \delta n,\;\; v^{\alpha} = 0 + v^{\alpha},$$

$$P^{\alpha} = P^{\alpha}_0 + \delta P^{\alpha}, R^{\alpha\beta} = 0 + \delta R^{\alpha\beta}. \qquad (13)$$

Substituting these relations into system of equations (3), (4), (11), (12) and (10) *and* neglecting nonlinear terms, we obtain a system of linear homogeneous equations in partial derivatives with constant coefficients. Passing to the following representation for small perturbations δf

$$\delta f = f(\omega,\mathbf{k})exp(-\imath\omega t + \imath\mathbf{k}\mathbf{r})$$

yields the homogeneous system of algebraic equations. The electric field strength is assumed to have a nonzero value. Expressing all the quantities entering the system of equations in terms of the electric field, we come to the equation

$$\Lambda\cdot E_z = 0,$$

where

$$\Lambda = \omega^2 - \frac{\hbar^2k^4}{4m^2} + \frac{\Upsilon k^2n_0}{2m} + 4\pi\sigma\frac{P_0^2k^2}{mn_0}$$

$$-\frac{2\pi\Upsilon k^4P_0^2}{m^2\omega^2 - \hbar^2k^4/4 + m\Upsilon k^2n_0 - m\Upsilon_2k^4n_0}.$$

In this case, the dispersion equation is

$$\Lambda = 0.$$

Solving this equation with respect to ω^2 we obtain a following results.

The dispersion characteristic for EE in BEC can be expressed in the form of

$$\omega^2 = \frac{1}{2m}\Biggl(-\frac{3}{2}\Upsilon n_0k^2 + \frac{\hbar^2k^4}{2m} + \Upsilon_2n_0k^4 + 4\pi\sigma\frac{P_0^2k^2}{n_0}$$

$$\pm\sqrt{\left(\frac{1}{2}\Upsilon n_0k^2 - \Upsilon_2n_0k^4 + 4\pi\sigma\frac{P_0^2k^2}{n_0}\right)^2 - 8\pi\Upsilon k^4P_0^2}\Biggr). \qquad (14)$$

In contrast to the non-polarized BEC, where only Bogoliubov's mode exists [6, 7, 14], a new wave solution appears in a polarized system due to the polarization dynamics. Bogoliubov's mode corresponds to a solution of the equation (14) with minus sign before the square root. New wave solution it is a wave of polarization. In general case presented by formula (14) the frequency of polarization wave in BEC depend on P_0 and Υ. To investigate the BEC polarization effect on the Bogoliubov's mode and the dispersion characteristic of the new solution we analyze extreme cases of the formula (14).

Formula (14) demonstrates that taking account of the BEC polarization dynamics leads to a new solution.

Let's start with the case when contributions of different term in formula (14) are comparable. In this case we studied dispersion numerically. For this purpose we represent formula (14) in the following form

$$\xi^2 = \Biggl(\frac{3}{4}\frac{a}{|a|} + \frac{k^2}{16\pi n_0|a|} - \frac{\theta a^3k^2}{16|a|} + \frac{6\pi\sigma a_{dd}}{|a|}$$

$$\pm\sqrt{\left(-\frac{1}{4}\frac{a}{|a|} + \frac{\theta a^3k^2}{16|a|} + \frac{6\pi\sigma a_{dd}}{|a|}\right)^2 + \frac{6\pi a_{dd}}{a}}\Biggr). \qquad (15)$$



FIG. 7. (Color online) Here we use $\lg k$, where k measured in cm^{-1}. The figure presents the dependence reduced frequency ξ on the wave vector k and DSL $a_{dd}/|a|$ for Bogoliubov's mode. Figure was obtained for repulsive SRI $a > 0$, $\Upsilon < 0$, $|a| = 10^{-7}$ cm, $n_0 = 10^{14}$ cm^{-3}, $\sigma = 1$, $\theta = 1$.



FIG. 8. (Color online) Here we use $\lg k$, where k measured in cm^{-1}. The figure presents the dependence reduced frequency ξ on the wave vector k for polarization wave at three different value of DSL a_{dd}: $a_{dd,1} = 0$ (red), $a_{dd,2} = |a|$ (green), $a_{dd,3} = 10|a|$ (blue), $a_{dd,4} = 100|a|$(black). Figure show increasing of ξ with increasing of a_{dd}. Figure obtained for repulsive SRI $a > 0$, $\Upsilon < 0$, and $|a| = 10^{-7}$ cm, $n_0 = 10^{14}$ cm^{-3}, $\sigma = 1$, $\theta = 1$.

where

$$\xi^2 \equiv \frac{\omega^2}{|\Upsilon| n_0 k^2/m}.$$

Here we use the length a_{dd} measures the strength of dipolar interaction, and defined by formula

$$a_{dd} = \frac{mP_0^2}{12\pi n_0^2 \hbar^2}.$$

Above we discuss that we do not consider the scattering process, but for convenience and analogy with the results of other paper we enter this parameter, although a_{dd} does not connect with scattering.



FIG. 9. (Color online) Here we use $\lg k$, where k measured in cm^{-1}. The figure shows behavior of reduced frequency ξ dependence on wave vector k for both solutions presented by formula (15). We can see the relative behavior of two solutions. Upper curve (blue) coincides to solution with sign ”+” in front of square root (polarization mode) and lower curve (red) correspond to the one with sign ”-” (Bogoliubov's mode). $a_{dd} = |a|$, $a > 0$, $\Upsilon < 0$ (repulsion), $|a| = 10^{-7}$ cm, $n_0 = 10^{14}$ cm^{-3}, $\sigma = 1$, $\theta = 1$. The figure presents comparison of the reduced frequencies dependence for Bogoliubov's and polarization modes. We can see that $\xi(k)$ for polarization mode lie above Bogoliubov's mode.



FIG. 10. (Color online) Here we use $\lg k$, where k measured in cm^{-1}. The figure presents the dependence reduced frequency ξ on the wave vector k for Bogoliubov's mode, $a < 0$, $\Upsilon > 0$ (attraction), DSL $a_{dd} = 0$ (Red, the right one), $a_{dd} = 0.0005|a|$ (Blue), $a_{dd} = 0.005|a|$ (Green), $a_{dd} = 100|a|$ (Red, the left one), $|a| = 10^{-7}$ cm, $n_0 = 10^{14}$ cm^{-3}, $\sigma = 1$, $\theta = 1$.

Here we describe the numerical analysis of formulas (15). Reduced frequency ξ for two waves presented by formula (15) exhibit on Fig.s 7 and 8. They are obtained for repulsive SRI ($a > 0$, $\Upsilon < 0$) at $a = 10^{-7}$cm and $n_0 = 10^{14}\text{cm}^{-3}$. These Fig.s show the reduced frequency $\xi(k)$ for different values of equilibrium polarization and, consequently, for the different DSL a_{dd}. Particularly, where the limit case a_{dd} is considered too.

Fig. 7 present solution with sign ”-” in front of square root in formula (14), this solution corresponds to the Bo-



FIG. 11. (Color online) Here we use $\lg k$, where k measured in cm^{-1}. The figure presents the dependence reduced frequency ξ on the wave vector k and DSL a_{dd} for polarization mode, $a < 0$, $\Upsilon > 0$ (attraction), $|a| = 10^{-7}$ cm, $n_0 = 10^{14}$ cm^{-3}, $\sigma = 1$, $\theta = 1$.



FIG. 12. (Color online) Here we use $\lg k$, where k measured in cm^{-1}. The figure presents the dependence reduced frequency ξ on the wave vector k for polarization mode, $a < 0$, $\Upsilon > 0$ (attraction), DSL $a_{dd} = 0$ (Red), $a_{dd} = |a|$ (Blue), $a_{dd} = 10|a|$ (Green), $a_{dd} = 100|a|$ (Brown), $|a| = 10^{-7}$ cm, $n_0 = 10^{14}$ cm^{-3}, $\sigma = 1$, $\theta = 1$.

goliubov's mode. For all $a_{dd}/|a|$ the $\xi(k)$ do not depend on $a_{dd}/|a|$.

Fig. 8 show strong dependence of reduced frequency of polarization mode ("+" in front of square root in (14)) on equilibrium polarization. On Fig.8 as on other Fig.s this dependence present via dependence on DSL divided by module of SL of SRI $a_{dd}/|a|$.

The form of $\xi(k)$ for two solutions has similar form. We present both solutions on Fig. 9 at fixed $a_{dd}/|a| = 1$ to compare. From Fig. 9 we see that reduced frequency of polarization mode is larger than one's for Bogoliubov's mode.

From the Fig.8 we obtained for polarization mode that polarization lead to growing of the reduced frequency ξ. At small wave vectors of order $10^4\mathrm{cm}^{-1}$ and for the $a_{dd} = |a|$ the reduced frequency ξ increases in 6 times. For the $a_{dd} = 100|a|$ one's increases in 60 times. With increasing of wave vector the influence of polarization became smaller. For example at $k = 10^6\mathrm{cm}^{-1}$ and $a_{dd} = 100|a|$ the reduced frequency ξ increase just in 1.6 times.



FIG. 13. (Color online) Here we use $\lg k$, where k measured in cm^{-1}. The figure presents the dependence reduced frequency ξ on the wave vector k for polarization mode. On this Fig. the region of instabilities is also shown. It is a region where a gap in curves. Figure obtained for attractive SRI ($a < 0$, $\Upsilon > 0$), $a_{dd} = 0$ (Red), $a_{dd} = 0.1|a|$ (Blue), $a_{dd} = 0.2|a|$ (Green), $a_{dd} = |a|$ (Black), $|a|$= 10^{-5} cm, $n_0 = 10^{14}$ cm^{-3}, $\sigma = 1$, $\theta = 1$.

For the case of attractive SRI in the absence of polarization there is instability of Bogoliubov's mode at small wave vectors. At the presence of small polarization the Bogoliubov's mode remains unstable, but area of stability become some wider as it shown on Fig10. There is fast widening of a region of stability at increasing of $a_{dd}/|a|$ Fig.11. Widening starts at approximately $a_{dd} = 0.02|a|$.

For the case of attractive SRI in the absence of polarization there is instability of polarization mode at small wave vector's (see Fig. 12). At $a_{dd} = 0$ we see instability of waves at wave vector k below $1.6 \cdot 10^4\mathrm{cm}^{-1}$. Existence of EDM leads to increasing of ξ and fast growing of the region of stability. Wave are stable up to $k = 10^3\mathrm{cm}^{-1}$ at $a_{dd}/|a| = 1$. Further increasing of $a_{dd}/|a|$ leads to increasing of ξ up to 64. ξ become equal to 64 at $a_{dd}/|a| \simeq 100$ and does not change at further increasing of $a_{dd}/|a|$.

Comparing Fig.s 10 and 12 we can see that reduced frequency ξ of polarization mode much larger than Bogoliubov's mode, as it was for repulsive SRI.

With increasing of strength of SRI the role of Υ_2 become more important. It's influence reveals at $|a| = 10^{-5}$cm for attractive SRI. In this case there is instability region at relatively large wave vectors. For described case the reduced frequency $\xi(k)$ is presented on Fig.s 13 and 14 at various $a_{dd}/|a|$. The region of instability is shown where. This instability appears because the expression under the square root in formula (14) became negative as it shown on Fig. 15.

Below, we analytically consider limit cases of general dispersion solution (14).

Here we consider the case when the effect of polarization



FIG. 14. (Color online) Here we use $\lg k$, where k measured in cm^{-1}. The figure presents the dependence reduced frequency ξ on the wave vector k for Bogoliubov's mode, $a<0$, $\Upsilon>0$ (attraction), DSL $a_{dd}=0$ (Red), $a_{dd}=0.1|a|$ (Black), $a_{dd}=0.2|a|$ (Blue), $a_{dd}=2|a|$ (Green), $|a|=10^{-5}$ cm, $n_0=10^{14}$ cm^{-3}, $\sigma=1$, $\theta=1$.



FIG. 15. (Color online) The figure presents the behavior of the square root in formula (31) (we conditionally designated this quantity as "sqrr") at the change of the wave vector k corresponding to the Fig. 10 at $a_{dd}=0.1|a|$, (square does not depend on concentration). $|a|=10^{-5}$ cm, $\sigma=1$, $\theta=1$. Here we use $\lg k$, where k measured in cm^{-1}.

is low compared to the contribution of short-range effects, i.e. when the members proportional to Υ and Υ_2 is comparably large. If so, then formula (14) takes the form

$$\omega_B^2=\frac{\hbar^2k^4}{4m^2}-\frac{\Upsilon n_0k^2}{m}+\frac{\Upsilon_2n_0k^4}{m}$$
$$-\frac{4\pi P_0^2k^2}{mn_0}\frac{\Upsilon(\sigma-1)-2\sigma\Upsilon_2k^2}{\Upsilon-2\Upsilon_2k^2} \qquad (16)$$

and

$$\omega_P^2=\frac{\hbar^2k^4}{4m^2}-\frac{1}{2}\frac{\Upsilon n_0k^2}{m}$$
$$+\frac{4\pi P_0^2k^2}{mn_0}\frac{\Upsilon(\sigma-1)-2\sigma\Upsilon_2k^2}{\Upsilon-2\Upsilon_2k^2}. \qquad (17)$$

Here we use indexes "B" for Bogoliubov's mode and "P" for new polarization mode. When deriving formulae (16) and (17) we expanded a sub-radical expression in (14) and took only first two terms of the expansion to estimate the influence of polarization on the wave dispersion.

As it follows from equations (11) and (12), changes in polarization can occur due to dipole-dipole interactions, SRI and quantum Bohm potential, i.e. to members proportional to $\hbar^2$. That's the reason for existence of other ways of polarization change when the contribution of equilibrium polarization P_0 is negligible. In the latter case, taken in a linear approximation, changes in the polarization do not affect the concentration evolution.

Formulae (14) are valid even in the absence of the external electrical field when equilibrium polarization equals zero $P_0=0$. Equations (14) in that case take the form

$$\omega_B^2=\frac{1}{m}\left(\frac{\hbar^2k^4}{4m}-\Upsilon n_0k^2+\Upsilon_2n_0k^4\right), \qquad (18)$$

$$\omega_P^2=\frac{1}{2m}\left(\frac{\hbar^2k^4}{2m}-\Upsilon n_0k^2\right). \qquad (19)$$

The waves of polarization could be existed at the absence of external electric field $E_0=0$. In this case the equilibrium state is no polarized and dipole direction of particles is distributed accidentally.

If the contribution of equilibrium polarization into BEC dispersion (14) is comparable to the contribution of SRI in the third order of the interaction radius Υ_2 relationships (14) are transformed into

$$\omega_B^2=\frac{\hbar^2k^4}{4m^2}-\frac{\Upsilon n_0k^2}{m}+\frac{\Upsilon_2n_0k^4}{m}+4\pi\frac{P_0^2k^2}{mn_0} \qquad (20)$$

and

$$\omega_P^2=\frac{\hbar^2k^4}{4m^2}-\frac{1}{2}\frac{\Upsilon n_0k^2}{m}+4\pi\frac{P_0^2k^2}{mn_0}(\sigma-1). \qquad (21)$$

Using Feshbach's resonance [71, 72] we can transform the SRI potential in such a way that $\Upsilon=0$ while $\Upsilon_2\neq0$. Formulas (14) in this situation turn into

$$\omega_B^2=\frac{\hbar^2k^4}{4m^2}+\frac{\Upsilon_2n_0k^4}{m} \qquad (22)$$

and

$$\omega_P^2=\frac{\hbar^2k^4}{4m^2}+4\pi\sigma\frac{P_0^2k^2}{mn_0}. \qquad (23)$$

In this paper we primarily focus on the influence of BEC polarization on its dispersion characteristics. So, let's consider the case when the contribution into the dispersion of the SRI at the first order of the interaction radius, i.e. terms proportional to Υ, is comparable to the contribution of polarization, and their total effect is much greater than the contribution of terms proportional to Υ_2. Here, (14) turns into

$$\omega_B^2 = \frac{\hbar^2 k^4}{4m^2} - \frac{\Upsilon n_0 k^2}{m}\biggl(\frac{(\sigma-1)}{\sigma}\biggr), \qquad (24)$$

and

$$\omega_P^2 = \frac{\hbar^2 k^4}{4m^2} + 4\pi\sigma\frac{P_0^2 k^2}{m n_0}$$

$$-\frac{1}{2}\frac{\Upsilon n_0 k^2}{m}\biggl(\frac{(\sigma+4)}{2\sigma}\biggr). \qquad (25)$$

polarization evolution at constant concentration

DBEC usually is the rare gas and we can not use condition incompressiblity, but we consider the case there concentration of particles is not change in time. We interested in dynamics of wave of polarization only. For this aim we need to use equations (11) and (12) at additional condition $n(\mathbf{r},t) = n_0 = const$. This is means the polarization change due to evolution of dipoles direction. Equation of polarization evolution (11) has no change in this case. In eq. (12) nonlinear term are disappeared and the term describing the SRI is simplified:

$$\partial_t R^{\alpha\beta}(\mathbf{r},t) + \partial^\gamma\Biggl(R^{\alpha\beta}(\mathbf{r},t)v^\gamma(\mathbf{r},t)$$

$$+R^{\alpha\gamma}(\mathbf{r},t)v^\beta(\mathbf{r},t) - P^\alpha(\mathbf{r},t)v^\beta(\mathbf{r},t)v^\gamma(\mathbf{r},t)\Biggr)$$

$$-\frac{\hbar^2}{4m^2}\partial_\beta\triangle P^\alpha(\mathbf{r},t) = \frac{n_0}{m}\Upsilon\partial^\beta\Biggl(P^\alpha(\mathbf{r},t)\Biggr)$$

$$+\frac{\sigma}{m n_0}P^\alpha(\mathbf{r},t)P^\gamma(\mathbf{r},t)\partial^\beta E^\gamma(\mathbf{r},t). \qquad (26)$$

and equation of field (10).

In this approximation exist only one wave solution exists, it is the wave of polarization. It's dispersion is

$$\omega^2 = \frac{\hbar^2 k^4}{4m^2} + \biggl(\frac{4\pi\sigma P_0^2}{m n_0} - \frac{n_0\Upsilon}{m}\biggr)k^2 \qquad (27)$$

In this case formula (27) reduces to

$$\omega^2 = \frac{4\pi\sigma P_0^2}{m n_0}.$$

For the particles with the EMD equal to 1 Debye at concentration 10^{12} cm^{-3} and wave vector $k \simeq 10^7$ cm^{-1} the frequency ω reach value of order $10^6 - 10^7$ s^{-1}.

Dispersion of 1D and 2D wave of polarization is described in Ref. [23].

IV. GENERATION OF WAVES IN POLARIZED BEC

In that section we consider the process of wave generation in BEC by means of the beam of neutral polarized particles. The interaction between beam and BEC has dipole-dipole origin.

To get the dispersion solution we use the system of QHD equations for each sort of particles (3), (4), (11), (12) and the equation of field (10). The equilibrium state of system is characterized by following values of the BEC parameters:

$$n = n_0 + \delta n,\ \ v^\alpha = 0 + v^\alpha,$$

$$P^\alpha = P_0^\alpha + \delta P^\alpha, R^{\alpha\beta} = 0 + \delta R^{\alpha\beta} \qquad (28)$$

and values of the beam parameters:

$$n_b = n_{0b} + \delta n_b,\ \ v_b^\alpha = U\delta^{z\alpha} + \delta v_b^\alpha,$$

$$P_b^\alpha = P_{0b}^\alpha + \delta P_b^\alpha,\ \ R_b^{\alpha\beta} = R_{0b}^{\alpha\beta} + \delta R_b^{\alpha\beta}. \qquad (29)$$

The polarization P_0^α is proportional to external electric field E_0^α. We consider the case then $\mathbf{E}_0 = [E_0 sin\varphi, 0, E_0 cos\varphi]$. In this case the tensor $R_{0b}^{\alpha\beta}$ has only two unequal to zero elements: $R_{0b}^{zx} = R_{0b} sin\varphi$ and $R_{0b}^{zz} = R_{0b} cos\varphi$. For the process, under consideration the dispersion relation is:

$$1 + 4\pi k^2\Biggl(\frac{P_0^2}{\omega^2 - \frac{\hbar^2 k^4}{4m^2} + \frac{\Upsilon n_0 k^2}{2m}}\times$$

$$\times\Biggl(\frac{\Upsilon k^2/(2m^2)}{\omega^2 - \frac{\hbar^2 k^4}{4m^2} + \frac{\Upsilon n_0 k^2}{m} - \frac{\Upsilon_2 n_0 k^4}{m}} - \frac{\sigma}{m n_0}\Biggr)$$

$$+\frac{1}{(\omega - k_z U)^2 - \frac{\hbar^2 k^4}{4m_b^2}}\times$$

$$\times\Biggl(\frac{2(\omega - k_z U)P_{0b}k_z(P_{0b}U - R_{0b})}{m_b n_{0b}\Bigl((\omega - k_z U)^2 - \frac{\hbar^2 k^4}{4m_b^2}\Bigr)} - \frac{\sigma_b P_{0b}^2}{m_b n_{0b}}\Biggr)\Biggr) = 0.$$

(30)

Using relation $P_{0b}U - R_{0b} = 0$, we can simplify the equation (30) and obtain

$$1 + \frac{\omega_D^2}{\omega^2 - \omega_1^2}\Biggl(\frac{\Upsilon n_0 k^2/(2m)}{\omega^2 - \omega_2^2} - \sigma\Biggr)$$

$$-\frac{\sigma_b\omega_{Db}^2}{(\omega - k_z U)^2 - \frac{\hbar^2 k^4}{4m_b^2}} = 0. \qquad (31)$$

In this formula the following designations are used

$$\omega_1^2 = \frac{\hbar^2 k^4}{4m^2} - \frac{\Upsilon n_0 k^2}{2m}, \qquad (32)$$

$$\omega_2^2 = \frac{\hbar^2 k^4}{4m^2} - \frac{\Upsilon n_0 k^2}{m} + \frac{\Upsilon_2 n_0 k^4}{m} \tag{33}$$

and

$$\omega_{Di}^2 = \frac{4\pi P_{0i}^2 k^2}{m_i n_{0i}}, \tag{34}$$

where i is the index of sorts of particles, the BEC or the beam.

The equation (31) has two beam related solutions, in the absence of BEC medium:

$$\omega = k_z U \pm \sqrt{\frac{\hbar^2 k^4}{4m_b^2} + \sigma_b \omega_{Db}^2}. \tag{35}$$

We will consider the possibilities of instabilities for the case of low-density beam, the limit case $\omega_{Db} \sim n_{0b} \to 0$. In this case we can neglect the last term in square root in (35). The resonance interaction beam with the BEC realizing if

$$k_z U \pm \frac{\hbar k^2}{2m_b} = \omega(k), \tag{36}$$

and could lead to instabilities. The quantity $\omega(k)$ is the dispersion of BEC modes (14). The frequency in this case can be presented in the form

$$\omega = k_z U \pm \frac{\hbar k^2}{2m_b} + \delta\omega. \tag{37}$$

Let us to consider two limit cases.

small frequency shift limit

In the limit case

$$\delta\omega \ll \hbar k^2/m \tag{38}$$

the frequency shift obtain in the form:

$$\delta\omega^2 = \pm\frac{2\sigma_b m_b m^2 \omega_{Db}^2 (\omega^2 - \omega_1^2)^2 (\omega^2 - \omega_2^2)^2}{\omega \omega_D^2 \hbar \Upsilon^2 n_0^2 k^6 W}, \tag{39}$$

where

$$W = 2\omega^2 - \omega_1^2 - \omega_2^2 - \frac{2m\sigma}{\Upsilon n_0 k^2}(\omega^2 - \omega_2^2)^2. \tag{40}$$

and the frequency ω determined with formula (14). The instabilities take place in the case $\delta\omega^2 < 0$. The sign of $\delta\omega^2$ is depend on the sign of W.

For the case resonance interaction of beam with the waves in BEC there are instabilities, for the first beam mode in (35) at $W < 0$ and for the second beam mode in formula (35) if $W > 0$. For the polarization mode W is positive. It means that the interaction of polarization mode with the second beam related mode results in the instability. For the Bogoliubov's mode the sign of W depend on σ.

We can consider the following cases:

(i) the contribution of equilibrium polarization to $\omega(k)$ is dominant; then $W > 0$;

(ii) the dominant contribution to $\omega(k)$ results from the term in $\omega(k)$ which is proportional to Υ, *and*, equilibrium polarization and SPI in TOIR give comparable contribution to $\omega(k)$. In this case there is

$$\sigma_0 = 1 + \frac{\Upsilon_2 n_0 k^4/m}{\Upsilon n_0 k^2/(2m) + 2\Upsilon_2 n_0 k^4/m - 4\omega_D^2}$$

. The sign of W varies at $\sigma = \sigma_0$. The dependence of W from σ is presented in the Table 1.

large frequency shift limit

In the limit case

$$\delta\omega \gg \hbar k^2/m \tag{41}$$

we have

$$\delta\omega = \xi \sqrt[3]{\frac{\sigma_b m^2 \omega_{Db}^2 (\omega^2 - \omega_1^2)^2 (\omega^2 - \omega_2^2)^2}{\omega \omega_D^2 \Upsilon^2 n_0^2 k^4 \mid W \mid}}, \tag{42}$$

where ξ equal to $\xi_1 = \sqrt[3]{1}$ for $W > 0$ or $\xi_{-1} = \sqrt[3]{-1}$ for $W < 0$. Evident forms of quantities ξ_1 and ξ_{-1} are

$$\xi_{-1} = [-1, \frac{-1 + \imath\sqrt{3}}{2}, \frac{-1 - \imath\sqrt{3}}{2}]$$

and

$$\xi_1 = [1, \frac{1 + \imath\sqrt{3}}{2}, \frac{1 - \imath\sqrt{3}}{2}].$$

The considerations concerning the sign of W, reflected in the table 1, are valid also for the limit condition (41).

From the formulas (39) and (42) we can see that external neutral particles beam leads to instabilities for both Bogoliubovs and polarization waves.

TABLE I. In this table the sign of the W are presented for the Bogoliubov's mode when SRI is prevailed.

	Υ>0 attraction	Υ<0 repulsion
$\sigma > \sigma_0$	+	-
$\sigma < \sigma_0$	-	+

V. CONCLUSION

In this work we developed the self-consistent method for description of dynamics of polarized BEC. This method accounts for the effect of polarization on changes in the concentration and in velocity field, which are determined in general by the continuity equation and Euler's equation. We derived the evolution equations of the polarization and the polarization current. The derived equations contain information about the influence of the interactions on the polarization evolution. We studied the effect of polarization on the BEC dynamics and the influence of SRI on the polarization evolution. An expression of SRI contribution in the equation of polarization current evolution via concentration, polarization and the SRI potential $\Upsilon = \Upsilon(U_{ij})$ was derived. With the assumption that the state of polarized Bose particles in the form of BEC can be described with some single-particle wave function. Changes in polarization due to SRI are shown to be determined at the first order of the interaction radius by the same interaction constant that occurs in Euler's equation and Gross-Pitaevskii equation.

The derivation of GP equation for polarized particles from QHD equation obtained in the paper. The condition of validity of GP equation was presented. Comparison of evolution equation of electrically polarized BEC and magnetized BEC was discussed. The differences between these equations are shown.

The physical meaning of self-consistent approximation was described. Suitability of self-consistent approximation for electrically polarized BEC was shown. Distinction of developed and used here self-consistent approximation for dipole-dipole interaction from scattering process are discussed.

Correct form of the dipole-dipole interaction Hamiltonian is discussed. The arguments for choosing of correct form of Hamiltonian for electric and magnetic dipoles are presented. Corresponding correct Hamiltonians were derived.

The dispersion of EE in the polarized BEC was analyzed. In the article, we show that polarization evolution in BEC causes a novel type of waves in BEC. The effect of polarization on the dispersion of the Bogoliubov's mode and the dispersion of a new wave mode were studied.

We show the possibility of wave generation in polarized BEC by means of monoenergetic beam of neutral polarized particles.

APPENDIX A

Hamiltonian's correct form

At derivation of system of QHD equations we need to write evident form of Hamiltonian of dipole-dipole interaction. In some works the dynamics of the magnetic dipole moment and of the EDM [32, 74] are analyzed in similar ways. Usual expressions of a Hamiltonian for dipole-dipole interaction are equal for electric and magnetic dipoles:

$$H_{dd}=\frac{\delta^{\alpha\beta}-3r^{\alpha}r^{\beta}/r^{2}}{r^{3}}d_{1}^{\alpha}d_{2}^{\beta}, \qquad (43)$$

In this paper for interaction of electric dipoles we used following Hamiltonian:

$$H_{dd}=-\partial^{\alpha}\partial^{\beta}\frac{1}{r}\cdot d_{1}^{\alpha}d_{2}^{\beta}.$$

There is well-known identity

$$-\partial^{\alpha}\partial^{\beta}\frac{1}{r}=\frac{\delta^{\alpha\beta}-3r^{\alpha}r^{\beta}/r^{2}}{r^{3}}+\frac{4\pi}{3}\delta^{\alpha\beta}\delta(\mathbf{r}), \qquad (44)$$

so we can see the difference between usually using Hamiltonian and one's used in this paper. The corrections of our selection followed from the fact that the equations obtained in the paper coincide to the Maxwell equations. At derivation of QHD equation in self-consistent approximation lead to field equation (10), we rewrite it here

$$\nabla\mathbf{E}(\mathbf{r},t)=-4\pi\nabla\mathbf{P}(\mathbf{r},t), \qquad (45)$$

but if we used Hamiltonian (43) we would obtain

$$\nabla\mathbf{E}(\mathbf{r},t)=\frac{8\pi}{3}\nabla\mathbf{P}(\mathbf{r},t), \qquad (46)$$

instead of (45).

It has been shown by Breit [76] that a Hamiltonian for spin-spin interaction, and, as a consequence, for the interaction of magnetic moments contains a term that is proportional to Dirac δ-function $\delta(r_1-r_2)d_1^{\alpha}d_2^{\beta}$ along with (43). The coefficient of the δ-function has been refined later [77] so that the Hamiltonian is in accord with Maxwell's free equations, such as $div\mathbf{B}=0$. The resultant expression for the spin-spin interaction Hamiltonian is:

$$H_{\mu\mu}=-\biggl(4\pi\delta_{\alpha\beta}\delta(\mathbf{r}_{12})+\nabla_1^{\alpha}\nabla_1^{\beta}(1/r_{12})\biggr)\mu_1^{\alpha}\mu_2^{\beta}, \qquad (47)$$

or

$$H_{\mu\mu}=\biggl(\frac{\delta^{\alpha\beta}-3r^{\alpha}r^{\beta}/r^{2}}{r^{3}}-\frac{8\pi}{3}\delta_{\alpha\beta}\delta(\mathbf{r})\biggr)\mu_1^{\alpha}\mu_2^{\beta}.$$

Actually, the δ functional term is used in literature [75], but authors followed to Breit and Landau [76], they used coefficient $-4\pi/3$ at δ function instead of $8\pi/3$, but this form of coefficient does not corresponds to Maxwell equation. It was shown in Ref. [77], where also were obtain correct coefficient presented in formula (47).

Thus, the differences in the dipole-dipole interactions of electric dipoles, and magnetic dipoles, must be taken into account in the development of theoretical field apparatus.

Method of equations derivation

The Schrödinger equation defines wave function in a 3N-dimensional configuration space. Physical processes in systems that involve large number of bodies occur in a three-dimensional physical space [78]. That's why a problem evolves of obtaining a quantum-mechanical description of a system of particles in terms of material fields e.g. concentration, momentum density, energy density and other fields of various tensor dimension that are defined in a three-dimension space.

The first step in MPQHD equation derivation is the definition of particles concentration. We define the quantum particles concentration as a quantum average of classic microscopic concentration in coordinate representation on many-particle wave function $\psi(R,t)$. Thus, the particles concentration evident form to be

$$n(\mathbf{r},t)=\int dR\sum_{i}\delta(\mathbf{r}-\mathbf{r}_{i})\psi^{*}(R,t)\psi(R,t),$$

where $dR=\prod_{i=1}^{N}d\mathbf{r}_{i}$. Differentiating this function over time and using the Schrodinger equation with the Hamiltonian (2), we derive the continuity equation (3). In the continuity equation the new quantity appears. We can differentiate this quantity over time and use the Schrodinger equation with the Hamiltonian (2). In the results we obtain equation for this quantity. In this way we can find an evolution equation for every additive physical quantity, but we mast know it's definition. The definition of new physical quantities appears at derivation of QHD equation. For example, the probability current definition appears at the derivation of continuity equation. At derivation of the current evolution equation (4) appears several quantities, one of them is the polarization presented below (48). Knowledge of the polarization definition give us ability to derive the equation of polarization evolution (12) and so on.

In equations (4), (10) polarization occurs in the form of

$$P^{\alpha}(\mathbf{r},t)=\int dR\sum_{i}\delta(\mathbf{r}-\mathbf{r}_{i})\psi^{*}(R,t)\hat{d}_{i}^{\alpha}\psi(R,t), \quad (48)$$

where $\mathbf{r}_{i}$ is the coordinate operator for i-th particle, $dR=\prod_{p=1}^{N}d\mathbf{r}_{p}$.

In right hand side of equation (12) where is the force-like field $F^{\alpha\beta}(\mathbf{r},t)$ which give rise to evolution of the polarization current $R^{\alpha\beta}(\mathbf{r},t)$. In general form, for $F^{\alpha\beta}(\mathbf{r},t)$, we can write:

$$F^{\alpha\beta}(\mathbf{r},t)=-\frac{1}{m}\partial_{\gamma}\Sigma^{\alpha\beta\gamma}(\mathbf{r},t)+\frac{1}{m}D^{\alpha\gamma}(\mathbf{r},t)\partial^{\beta}E^{\gamma}(\mathbf{r},t). \quad (49)$$

A tensor

$$D^{\alpha\beta}(\mathbf{r},t)=\int dR\sum_{i}\delta(\mathbf{r}-\mathbf{r}_{i})d_{i}^{\alpha}d_{i}^{\beta}\psi^{*}(R,t)\psi(R,t), \quad (50)$$

occurs in the term that represents a dipole-dipole interaction and an interaction of the dipole with external electrical field. Based on the reasons of dimensions this value can be approximately presented as

$$D^{\alpha\beta}(\mathbf{r},t)=\sigma\frac{P^{\alpha}(\mathbf{r},t)P^{\beta}(\mathbf{r},t)}{n(\mathbf{r},t)}. \quad (51)$$

A SRI causes the tensor $\Sigma^{\alpha\beta\gamma}(\mathbf{r},t)$ to occur in the equation (12). Taken at the first order of the interaction radius it has the form

$$\Sigma^{\alpha\beta\gamma}(\mathbf{r},t)=-\frac{1}{2}\int dR\sum_{i,j\neq i}\delta(\mathbf{r}-\mathbf{R}_{ij})$$

$$\times\frac{r_{ij}^{\beta}r_{ij}^{\gamma}}{r_{ij}}\frac{\partial U_{ij}}{\partial r_{ij}}\psi^{*}(R,t)\hat{d}_{i}^{\alpha}\psi(R,t). \quad (52)$$

Tensor $\Sigma^{\alpha\beta\gamma}(\mathbf{r},t)$ describe the influence of SRI on evolution of polarization. Eq. (52) describe the SRI.

If we apply the procedure described in [14] to the calculation of the quantum stress tensor, and neglect the contribution of thermal excitations, $\Sigma^{\alpha\beta\gamma}(\mathbf{r},t)$ for BEC takes a form of

$$\Sigma_{BEC}^{\alpha\beta\gamma}(\mathbf{r},t)=-\frac{1}{2}\Upsilon\delta^{\beta\gamma}n(\mathbf{r},t)P^{\alpha}(\mathbf{r},t). \quad (53)$$

Formula (53) is obtained for the case where particles located in state with the lowest energy, which could be described by one particle wave function. This state may be the product of strong interaction.

Tensor $\Sigma^{\alpha\beta\gamma}(\mathbf{r},t)$ is, therefore, like the quantum stress tensor $\sigma^{\alpha\beta}(\mathbf{r},t)$ in the momentum balance equation (4), dependent on Υ at the first order of the interaction radius [14].

APPENDIX B: DERIVATION OF THE NON-LINEAR SCHRÖDINGER EQUATION

In Ref. [14] the derivation of GP equation was performed for non-polarized particles from QHD equation, here we present one for polarized particles. The NLSE comes about from the continuity equation and the Cauchy integral of the momentum balance equation. The Cauchy integral exists provided that the velocity field has the form

$$v^{\alpha}(\mathbf{r},t)=\frac{1}{m}\partial^{\alpha}\theta(\mathbf{r},t) \quad (54)$$

where $\theta(\mathbf{r},t)$ is a velocity field potential.

Starting from QHD equations we can derive an equation for evolution of the model function defined in terms of hydrodynamic variables. Thus a macroscopic single-particle wave function may be defined as

$$\Phi(\mathbf{r},t)=\sqrt{n(\mathbf{r},t)}\exp\biggl(\frac{\imath}{\hbar}m\theta(\mathbf{r},t)\biggr). \quad (55)$$

If we differentiate this function with respect to time and apply QHD equations then given

$$\imath\hbar\partial_t\Phi(\mathbf{r},t)=\Biggl(-\frac{\hbar^2}{2m}\nabla^2+\Gamma(\mathbf{r},t)+g\mid\Phi(\mathbf{r},t)\mid^2$$

$$-\int_{\mathbf{r}_0}^{\mathbf{r}}d\mathbf{r}'\frac{1}{n(\mathbf{r}',t)}P^\beta(\mathbf{r}',t)\times$$

$$\times\nabla'\int d\mathbf{r}''G^{\beta\gamma}(\mathbf{r}',\mathbf{r}'')P^\gamma(\mathbf{r}'',t)\Biggr)\Phi(\mathbf{r},t),\qquad(56)$$

here we use designation $\Gamma(\mathbf{r},t)$ described by formula

$$\Gamma(\mathbf{r},t)=\int_{\mathbf{r}_0}^{\mathbf{r}}d\mathbf{r}'\frac{\nabla p(\mathbf{r}',t)}{n(\mathbf{r}',t)}.\qquad(57)$$

The quantity $\Gamma(\mathbf{r},t)$ can be considered as chemical potential $\mu(\mathbf{r},t)$ and we used this fact below. The equation obtained (56) has the form of NLSE. NLSE with DDI were obtained on Ref. [23], where the sign ”-” was lost before integral in formulas (A6) and (A8)

NLSE (56) describes collective characteristics in a system of many particles. This follows from the derivation of NLSE and from the definition of a many-particle wave function $\Phi(\mathbf{r},t)$ (55). Function $\Gamma(\mathbf{r},t)$ (57) is the contribution of the kinetic pressure and does not contain any interaction.

If we introduce $d^\alpha(\mathbf{r},t)$ as $d^\alpha(\mathbf{r},t)=P^\alpha(\mathbf{r},t)/n(\mathbf{r},t)$ and if also we suppose that $d^\alpha(\mathbf{r},t)$ is a constant d^α from eq. (56) we obtain

$$\imath\hbar\partial_t\Phi(\mathbf{r},t)=\Biggl(-\frac{\hbar^2}{2m}\nabla^2+\mu(\mathbf{r},t)+g\mid\Phi(\mathbf{r},t)\mid^2$$

$$-d^\beta d^\gamma\int d\mathbf{r}'G^{\beta\gamma}(\mathbf{r},\mathbf{r}')|\Phi(\mathbf{r}',t)|^2\Biggr)\Phi(\mathbf{r},t),\qquad(58)$$

where we used $n(\mathbf{r},t)=|\Phi(\mathbf{r},t)|^2$.

At temperature T equal to zero in external electric field $\mathbf{E}_0$ all dipoles directed parallel to external field. If $\mathbf{E}_0=E_0\mathbf{e}_z$ we have $\mathbf{d}=d_0\mathbf{e}_z$. In this case in equation (58) only one component of Green function $G^{\beta\gamma}$ of DDI is: $G^{zz}=(1-3z^2/r^2)/r^3+4\pi/3\cdot\delta(\mathbf{r})$. As $z^2/r^2=\cos^2\theta$ and in the absence of δ function from (58) we obtain the GP equation (1) presented in introduction.

The GP equation was obtained for the case of system of parallel dipoles. We suppose it can be approximately used for analyze of dipoles whose direction slowly changed in space.

APPENDIX C

Here we consider QHD equations for the spinning particles and it’s difference from the one’s for electrically polarized.

In Euler equation contribution of magnetization **M** appear instead of polarization **P**. Method of QHD also allow to obtain equation of evolution of magnetization to be

$$\partial_t M^\alpha(\mathbf{r},t)+\nabla^\beta J_M^{\alpha\beta}(\mathbf{r},t)$$

$$=\frac{\gamma}{\hbar}\varepsilon^{\alpha\beta\gamma}\Biggl(M^\beta(\mathbf{r},t)B^\beta(\mathbf{r},t)\Biggr).\qquad(59)$$

where $J_M^{\alpha\beta}$ arises. Vanishing by thermal motion we have $J_M^{\alpha\beta}=M^\alpha v^\beta$.

For obtaining of QHD equations we started from many-particle Schrodinger equation

$$\imath\hbar\partial_t\psi_s(R,t)=\Biggl(\Biggl(\sum_i\Biggl(\frac{p_i^2}{2m_i}-\gamma_i\hat{s}_i^\alpha B^\alpha_{i(ext)}\Biggr)$$

$$+\frac{1}{2}\sum_{i,j\neq i}\Biggl(U_{ij}-\gamma_i\gamma_j G_{ij}^{\alpha\beta}\hat{s}_i^\alpha\hat{s}_j^\beta\Biggr)\Biggr)\psi\Biggr)_s(R,t).\qquad(60)$$

where we include short range and spin-spin interactions, and action of external magnetic field on spin. In Schrodinger equation (60) we used following designations: γ_i is the gyromagnetic ratio, $\mathbf{p}_i=-\imath\hbar\nabla_i$ is the operator of momentum, U_{ij} present short range interaction, the Green function of spin-spin interaction has form $G_{ij}^{\alpha\beta}=4\pi\delta_{\alpha\beta}\delta(\mathbf{r}_{12})+\nabla_1^\alpha\nabla_1^\beta(1/r_{12})$.

For spin matrixes $\hat{s}_i^\alpha$ the commutation relations are

$$[\hat{s}_i^\alpha,\hat{s}_j^\beta]=\imath\delta_{ij}\varepsilon^{\alpha\beta\gamma}\hat{s}_i^\gamma.$$

Thereby we consider Bose particles we present here the evident form of the spin matrixes $\hat{s}_i^\alpha$ for particles with spin equal to 1:

$$\hat{s}_x=\tfrac{1}{\sqrt{2}}\begin{pmatrix}0&1&0\\1&0&1\\0&1&0\end{pmatrix},\ \hat{s}_y=\tfrac{1}{\sqrt{2}}\begin{pmatrix}0&-\imath&0\\\imath&0&-\imath\\0&\imath&0\end{pmatrix},$$

$$\hat{s}_z=\begin{pmatrix}1&0&0\\0&0&0\\0&0&-1\end{pmatrix}.$$

For derivation of QHD equation from Schrodinger equation we start as usual from definition of concentration of particles around point **r** of physical space:

$$n(\mathbf{r},t)=\sum_s\int dR\sum_i\delta(\mathbf{r}-\mathbf{r}_i)\psi_s^+(R,t)\psi_s(R,t),$$

where $dR=\prod_{i=1}^{N}d\mathbf{r}_i$. In this case we obtain an equations analogous to (3) and (4) where instead polarization $\mathbf{P}(\mathbf{r},t)$ appear magnetization $\mathbf{M}(\mathbf{r},t)$.

The magnetization arise in the form

$$M^{\alpha}(\mathbf{r},t)=\sum_{s}\int dR\sum_{i}\delta(\mathbf{r}-\mathbf{r}_i)\times$$

$$\times\,\gamma_i\psi^{+}(R,t)_s(\hat{s}^{\alpha}_i\psi(R,t))_s. \qquad (61)$$

Differentiating magnetization (61) with respect to time and using Schrodinger equation (60) we came to equation (59).

More details for obtaining of QHD equations for spinning particles are presented in Ref.s [23], [69].

We can perform derivation of NLSE for spinning particles, in the result we have obtained an equation analogous to (56). In the case parallel dipoles we have following equation, where the difference between Green function of spin-spin EDM interactions is accounted

$$\imath\hbar\partial_t\Phi(\mathbf{r},t)=\Biggl(-\frac{\hbar^2}{2m}\nabla^2+\mu(\mathbf{r},t)+g\mid\Phi(\mathbf{r},t)\mid^2$$

$$+\mu_0^2\int d\mathbf{r}'\frac{1-3\cos^2\theta'}{|\mathbf{r}-\mathbf{r}'|^3}|\Phi(\mathbf{r}',t)|^2-\frac{8\pi}{3}\mu_0^2|\Phi(\mathbf{r},t)|^2\Biggr)\Phi(\mathbf{r},t). \qquad (62)$$

Thereby, feature of spin-spin interaction give us additional term in GP equation for spinning particles. This term caused by long-range interaction, but it has form analogous to SRI.

---

* andreevpa@physics.msu.ru
† lsk@phys.msu.ru

[1] Patrick Koberle, Holger Cartarius, Tomaz Fabcic, Jorg Main and Gunter Wunner, New Journal of Physics **11**, 023017 (2009).
[2] Lincoln D. Carr, David DeMille, Roman V. Krems and Jun Ye, New Journal of Physics **11**, 055049 (2009).
[3] Thierry Giamarchi, Christian Ruegg and Oleg Tchernyshyov, Nature Phys. **4**, 198 (2008).
[4] K.-K. Ni, S. Ospelkaus, D. J. Nesbitt, J. Ye and D. S. Jin, Phys. Chem. Chem. Phys. **11**, 9626 (2009).
[5] A. Griffin, *Excitations in a Bose-Condensed Liquid* (Cambridge University Press, New York, 1993).
[6] N. Bogoliubov, J. Phys. (Moscow) **11**, 23 (1947).
[7] F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari, Rev. Mod. Phys. **71**, 463 (1999).
[8] H. Pu, W. Zhang, M. Wilkens, and P. Meystre, Phys. Rev. Lett. **88**, 070408 (2002).
[9] J. Steinhauer, R. Ozeri, N. Katz, and N. Davidson, Phys. Rev. Lett. **88**, 120407 (2002).
[10] Chiara Menotti, S. Stringari, Phys. Rev. A **66**, 043610 (2002).
[11] A. Banerjee, M. P. Singh, Phys. Rev. A **66**, 043609 (2002).
[12] H. Shibata, N. Yokoshi, S. Kurihara, Phys. Rev. A **75**, 053615 (2007).
[13] M. Gupta and K. Rai Dastidar, Phys. Rev. A **81**, 063631 (2010).
[14] P. A. Andreev, L. S. Kuz'menkov, Phys. Rev. A **78**, 053624 (2008).
[15] S. Stringari, Phys. Rev. Lett. **77**, 2360 (1996).
[16] G. M. Falco, A. Pelster, and R. Graham, Phys. Rev. A **76**, 013624 (2007).
[17] A. Banerjee, Phys. Rev. A **76**, 023611 (2007); A. Banerjee, J. Phys. B **42**, 235301 (2009).
[18] L. Santos, G. V. Shlyapnikov, and M. Lewenstein, Phys. Rev. Lett. **90**, 250403 (2003).
[19] D. H. J. ODell, S. Giovanazzi, and G. Kurizki, Phys. Rev. Lett. **90**, 110402 (2003).
[20] S. Giovanazzi and D. H. J. ODell, Eur. Phys. J. D **31**, 439 (2004).
[21] Uwe R. Fischer, Phys. Rev. A **73**, 031602(R) (2006).
[22] C. Ticknor, R. M. Wilson, and J. L. Bohn, Phys. Rev. Lett. **106**, 065301 (2011).
[23] P. A. Andreev, L. S. Kuzmenkov and M. I. Trukhanova, Phys. Rev. B **84**, 245401 (2011).
[24] Qiuzi Li, E. H. Hwang, and S. Das Sarma, Phys. Rev. B **82**, 235126 (2011).
[25] S. Yi and L. You, Phys. Rev. A, **61**, 041604(R) (2000).
[26] K. Goral, K. Rzazewski, and T. Pfau, Phys. Rev. A **61**, 051601(R) (2000).
[27] L. Santos, G.V. Shlyapnikov, P. Zoller, and M. Lewenstein, Phys. Rev. Lett. **85**, 1791 (2000).
[28] S. Yi and L. You, Phys. Rev. A, **63**, 053607 (2001).
[29] K. Goral and L. Santos, Phys. Rev. A **66**, 023613 (2002).
[30] P. Szankowski, M. Trippenbach, E. Infeld, Ge. Rowlands, Phys. Rev. Lett. **105**, 125302 (2010).
[31] R. W. Cherng and E. Demler, Phys. Rev. Lett. **103**, 185301 (2009).
[32] T. Lahaye, C. Menotti, L. Santos, M. Lewenstein and T. Pfau, Rep. Prog. Phys. **72**, 126401 (2009).
[33] Yongyong Cai, Matthias Rosenkranz, Zhen Lei, and Weizhu Bao, Phys. Rev. A **82**, 043623 (2010).
[34] I. Sapina, T. Dahm, and N. Schopohl, Phys. Rev. A **82**, 053620 (2010).
[35] R. M. W. van Bijnen, N. G. Parker, S. J. J. M. F. Kokkelmans, A. M. Martin, and D. H. J. O'Dell, Phys. Rev. A **82**, 033612 (2010).
[36] S. Komineas and N. R. Cooper, Phys. Rev. A **75**, 023623 (2007).
[37] R. M. Wilson, S. Ronen, and J. L. Bohn, Phys. Rev. Lett. **104**, 094501 (2010).
[38] YuanYao Lin, Ray-Kuang Lee, Yee-Mou Kao, and Tsin-Fu Jiang, Phys. Rev. A **78**, 023629 (2008).
[39] T. F. Jiang and W. C. Su, Phys. Rev. A **74**, 063602 (2006).
[40] G. Gligoric, A. Maluckov, M. Stepic, L. Hadzievski, and B. A. Malomed, Phys. Rev. A **81**, 013633 (2010).
[41] T. Lahaye, T. Koch, B. Frohlich, M. Fattori, J. Metz, A. Griesmaier, S. Giovanazzi, T. Pfau, Nature **448**, 672 (2007).
[42] Y. Yamaguchi, T. Sogo, T. Ito, and T. Miyakawa, Phys. Rev. A **82**, 013643 (2010).
[43] Zhen-Kai Lu, G. V. Shlyapnikov, arXiv:1111.7114.
[44] Anne-Louise Gadsbolle, and G. M. Bruun, arXiv:1112.2846.
[45] L. S. Kuz'menkov and S. G. Maksimov, Teor. i Mat. Fiz., **118**, 287 (1999) [Theoretical and Mathematical Physics **118**, 227 (1999)].

[46] P. A. Andreev, arXiv: 1201.0779.
[47] P. A. Andreev, arXiv:1109.0896.
[48] Daw-Wei Wang, New Journal of Physics **10**, 053005 (2008).
[49] L. P. Pitaevskii, E.M. Lifshitz ”Physical Kinetics” (kinetic theory), Vol. 10 of Course of Theoretical Physics (Pergamon, London, 1981).
[50] A. A. Vlasov, J. Exp. Theor. Phys. **8**, 291 (1938); A. A. Vlasov Sov. Phys. Usp. **10**, 721 (1968).
[51] A. R. P. Lima, A. Pelster, Phys. Rev. A **81**, 063629 (2010).
[52] H. Gomi, T. Imai, A. Takahashi, and M. Aihara, Phys. Rev. B **82**, 035101 (2010).
[53] G. Quemener and J. L. Bohn, Phys. Rev. A **83**, 012705 (2011).
[54] R. M. Lutchyn, E. Rossi, and S. Das Sarma, Phys. Rev. A **82**, 061604(R) (2010).
[55] R. Liao and J. Brand, Phys. Rev. A **82**, 063624 (2010).
[56] D. Baillie and P. B. Blakie, Phys. Rev. A **82**, 033605 (2010).
[57] K. Gillen-Christandl and B. D. Copsey, Phys. Rev. A **83**, 023408 (2011).
[58] H. Deng, H. Haug, Y. Yamamoto, Rev. Mod. Phys. **82**, 1489 (2010).
[59] H. Haug, and A. Jauho, *Quantum Kinetics in Transport and Optics of Semiconductors*, 2nd ed. (Springer, Berlin, 2008).
[60] T. Kuhn, *Theory of Transport Properties of Semiconductor Nanostructures*, edited by E. Scholl (Springer, New York, 1997).
[61] U. Vogl, M. Weitz, Nature **461**, 70 (2009).
[62] M. Sheik-Bahae, D. Seletskiy, Nature Photonics **3**, 680 (2009).
[63] N. N. Rosanov, A. G. Vladimirov, D. V. Skryabin, W. J. Firth, Phys. Lett. A. **293**, 45 (2002).
[64] E. Braaten, H.-W. Hammer, and Shawn Hermans, Phys. Rev. A. **63**, 063609 (2001).
[65] E. Braaten, L. Platter, Phys. Rev. Lett. **100**, 205301 (2008).
[66] P. A. Andreev, L. S. Kuzmenkov, arXiv:1105.5537.
[67] A. Bret, M.-C. Firpo, and C. Deutsch, Phys. Rev. E **70**, 046401 (2004).
[68] P. A. Andreev, L.S. Kuz’menkov, Physics of Atomic Nuclei **71**, N.10, 1724 (2008); P. A. Andreev and L. S. Kuz’menkov, PIERS Proceedings, p. 1047, March 20-23, Marrakesh, MOROCCO 2011.
[69] P. A. Andreev and L. S. Kuzmenkov, arXiv:1109.3335.
[70] P. A. Andreev and L. S. Kuzmenkov, arXiv:1106.0822
[71] Cheng Chin, R. Grimm, P. Julienne and E. Tiesinga, Rev. Mod. Phys. **82**, 1225 (2010).
[72] I. Bloch, J. Dalibard, W. Zwerger, Rev. Mod. Phys. **80**, 885 (2008).
[73] I. A. Akhiezer, *Plasma electrodynamics*, (Pergamon Press, 1975).
[74] S. Yi and L. You, Phys. Rev. A **66**, 013607 (2002).
[75] Claudia Eberlein, Stefano Giovanazzi, and Duncan H. J. O’Dell, Phys. Rev. A **71**, 033618 (2005).
[76] G. Breit, Phys. Rev. **34**, 553 (1929); V.B. Berestetskii, E.M. Lifshitz, L.P. Pitaevskii (1982). Quantum Electrodynamics. Vol. 4 (2nd ed.). Butterworth-Heinemann; V. Yu. Lazur, S. I. Myhalyna, and O. K. Reity, Phys. Rev. A **81**, 062707 (2010).
[77] L. S. Kuz’menkov, S. G. Maksimov, and V. V. Fedoseev, Theor. Math. Fiz. **126**, 136 (2001) [Theoretical and Mathematical Physics **126**, 110 (2001)].
[78] S. Goldstein, Physics Today. **51**, N. 3, 42 (1998); S. Goldstein, Physics Today. **51**, N. 4, 38 (1998).